



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Book
Search	PubMed	<input checked="" type="checkbox"/> for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		

Display	Abstract	<input checked="" type="checkbox"/>	Show	20	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Send to	Text	<input type="checkbox"/>
---------	----------	-------------------------------------	------	----	--------------------------	------	--------------------------	---------	------	--------------------------

Entrez PubMed

☐ 1: Pharmacol Rev 2002 Sep;54(3):375-429

Related Articles, Links

Full text article at
www.pharmrev.org

The therapeutic potential of poly(ADP-ribose) polymerase inhibitors.

Virag L, Szabo C.

Inotek Pharmaceutical Corp., Beverly, Massachusetts 01915, USA.

Poly(ADP-ribose) polymerase-1 (PARP-1) is a member of the PARP enzyme family consisting of PARP-1 and several recently identified novel poly(ADP-ribosylating) enzymes. PARP-1 is an abundant nuclear protein functioning as a DNA nick-sensor enzyme. Upon binding to DNA breaks, activated PARP cleaves NAD(+) into nicotinamide and ADP-ribose and polymerizes the latter onto nuclear acceptor proteins including histones, transcription factors, and PARP itself. Poly(ADP-ribosylation) contributes to DNA repair and to the maintenance of genomic stability. On the other hand, oxidative stress-induced overactivation of PARP consumes NAD(+) and consequently ATP, culminating in cell dysfunction or necrosis. This cellular suicide mechanism has been implicated in the pathomechanism of stroke, myocardial ischemia, diabetes, diabetes-associated cardiovascular dysfunction, shock, traumatic central nervous system injury, arthritis, colitis, allergic encephalomyelitis, and various other forms of inflammation. PARP has also been shown to associate with and regulate the function of several transcription factors. Of special interest is the enhancement by PARP of nuclear factor kappa B-mediated transcription, which plays a central role in the expression of inflammatory cytokines, chemokines, adhesion molecules, and inflammatory mediators. Herein we review the double-edged sword roles of PARP in DNA damage signaling and cell death and summarize the underlying mechanisms of the anti-inflammatory effects of PARP inhibitors. Moreover, we discuss the potential use of PARP inhibitors as anticancer agents, radiosensitizers, and antiviral agents.


Publication Types:

- Review
- Review, Academic

PMID: 12223530 [PubMed - indexed for MEDLINE]

PubMed Services

Related Resources



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Book
Search	PubMed	<input checked="" type="checkbox"/>	for		Go	Clear		
Limits		Preview/Index		History		Clipboard		Details
Display	Abstract	<input checked="" type="checkbox"/>	Show:	20	Sort	<input checked="" type="checkbox"/>	Send to	Text

Entrez PubMed

☐ 1: Curr Pharm Biotechnol 2002 Sep;3(3):275-83[Related Articles, Links](#)

PubMed Services

Modulating poly (ADP-ribose) polymerase activity: potential for the prevention and therapy of pathogenic situations involving DNA damage and oxidative stress.

Decker P, Muller S.

Institute for Cell Biology, Department of Immunology; Auf der Morgenstelle, Tübingen, Germany. patrice.decker@uni-tuebingen.de

Related Resources

Poly (ADP-ribose) polymerase is a zinc-finger DNA-binding enzyme which detects and signals DNA strand breaks generated either directly during base excision repair, or indirectly by genotoxic agents such as oxygen radicals. In response to genotoxic injury, PARP catalyses the synthesis of poly (ADP-ribose), from its substrate beta-NAD⁺ and this polymer is covalently attached to several nuclear proteins and PARP itself. As a result, PARP converts DNA breaks into intracellular signals which activate DNA repair programs or cell death options. Several studies have also shown that PARP is involved in either necrosis and subsequent inflammation or apoptosis. Although this enzyme is not indispensable during the latter cell death program, it has been demonstrated that PARP plays a facilitating role in this process. PARP is activated at an intermediate stage of apoptosis and is then cleaved and inactivated at a late stage by apoptotic proteases, namely caspase-3/CPP-32/Yama/apopain and caspase-7. This cleavage prevents necrosis during apoptosis, avoiding inflammation. All these functions, and the observation that PARP is an abundant and highly conserved enzyme, suggest that this enzyme plays a pivotal role, particularly in the maintenance of genomic DNA stability, apoptosis and in the response to oxidative stress. Since these situations are found in cancer, inflammation, autoimmunity (such as diabetes), myocardial dysfunction, certain infections, ageing and radiation/chemical exposure, attempts have been made to modulate PARP activity. With regard to the increasing interest towards PARP, the aim of this review is to explain the cellular role of PARP and the advantages of modulating its activity in diverse preventive or therapeutic strategies.

Publication Types:

- Review
- Review, Tutorial

**Mutation Research/DNA Repair**

Volume 460, Issue 1, 30 June 2000, Pages 1-15

doi:10.1016/S0921-8777(00)00016-1 Cite or link using doi
Copyright © 2000 Elsevier Science B.V. All rights reserved.

This Document► **Abstract**

- [Full Text + Links](#)
- [PDF \(255 K\)](#)

Review**Actions**

- [E-mail Article](#)

Poly(ADP-ribose) polymerase-1: what have we learned from the deficient mouse model?

Sydney Shall¹, and Gilbert de Murcia,

UPR 9003 du Centre National de la Recherche Scientifique, Laboratoire conventionné avec le Commissariat à l'Energie Atomique, Ecole Supérieure de Biotechnologie de Strasbourg, boulevard Sébastien Brant, F-67400 Illkirch-Graffenstaden, France


Received 8 February 2000; revised 30 March 2000; accepted 5 April 2000. Available online 13 June 2000.

Abstract

Poly (ADP-ribose) polymerase (113 kDa; PARP-1) is a constitutive factor of the DNA damage surveillance network developed by the eukaryotic cell to cope with the numerous environmental and endogenous genotoxic agents. This enzyme recognizes and is activated by DNA strand breaks. This original property plays an essential role in the protection and processing of the DNA ends as they arise in DNA damage that triggers the base excision repair (BER) pathway. The generation, by homologous recombination, of three independent deficient mouse models have confirmed the caretaker function of PARP-1 in mammalian cells under genotoxic stress. Unexpectedly, the knockout strategy has revealed the instrumental role of PARP-1 in cell death after ischemia–reperfusion injury and in various inflammation process. Moreover, the residual PARP activity found in PARP-1 deficient cells has been recently attributed to a novel DNA damage-dependent poly ADP-ribose polymerase (62 kDa; PARP-2), another member of the expanding PARP family that, on the whole, appears to be involved in the genome protection. The present review summarizes the recent data obtained with the three PARP knockout mice in comparison with the chemical inhibitor approach.

Author Keywords: Cellular response to DNA damage; Genomic instability; Base excision repair; Type I Diabetes; NF- κ B; NAD⁺ metabolism

¹ Current address: Department of Molecular Medicine, The Rayne Institute, 123 Coldharbour Lane, King's College School of Medicine and Dentistry, London, SE5 9NU, England.

 Corresponding author. Tel.: +33-3-88-65-53-68; fax: +33-3-88-65-53-43; email: demurcia@esbs.u-strasbg.fr

Mutation Research/DNA Repair

Volume 460, Issue 1 , 30 June 2000 , Pages 1-15

This Document

► **Abstract**

- [Full Text + Links](#)
- [PDF \(255 K\)](#)

Actions

- [E-mail Article](#)

[Home](#)

[Journals](#)

[Abstract Databases](#)

[Reference Works](#)

[My Alerts](#)

[My Profile](#)

[? Help](#)

Send [feedback](#) to ScienceDirect

Software and compilation © 2003 ScienceDirect. All rights reserved.

ScienceDirect® is an Elsevier Science B.V. registered trademark.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.